

Viridans Just-a-Minute : Tutorial 1

Task:

Investigate Alpine Egernia (*Egernia guthega*).


- Find the animal in Just-a-Minute Animals.
- View full-sized photographs of the species
- Print a combined photograph, written description and distribution map of the species.
- Plot the distribution of the species on a map of Victoria.
- Display the same distribution on a map high resolution map centred on the Victorian Alps.
- Export that map as a Windows Metafile (WMF) document.

The products of this task should be:

- A printed page with a photograph, written description and distribution map of Alpine Egernia
- A printed page or saved WMF document with the distribution of Alpine Egernia plotted as one-minute grids on a 1:100,000 map of Victorian Alps.

Just a Minute (Standard)
Just a Minute (1:1,000,000 to 1:100,000)
Just a Minute (1:1,000,000 to 1:100,000 - Low Contrast)
Help
Close

Select this option



Full Species List

Area Species List

Search an Area

Help

Close

Select this option

Just-a-Minute
Victorian Animals
2012



Just a Minute Victorian Animals - 1:1,000,000 to 1:100,000

Enter Common Name

alp eger

Find Name

Sort on Scientific Name

Common Name Order : Full List : 780 species

Click on column buttons for interpretation of this list

F	A	S	O	Common Name	Scientific Name	Family Name	Last Date
				Adelie Penguin	<i>Pygoscelis adeliae</i>	Spheniscidae	No Records
				Agile Antechinus	<i>Antechinus agilis</i>	Dasyuridae	2000-2010
f		e		Alpine Bog Skink	<i>Pseudemoia cryodroma</i>	Scincidae	2000-2010
f		c		Alpine Egernia	<i>Egernia guthega</i>	Scincidae	2000-2010
f		e		Alpine She-oak Skink	<i>Cyclodomorphus praealtus</i>	Scincidae	2000-2010
f	V	c		Alpine Tree Frog	<i>Litoria verreauxii alpina</i>	Hylidae	2000-2010
f		c		Alpine Water Skink	<i>Eulamprus kosciuskoi</i>	Scincidae	2000-2010
				Andrew's Beaked Whale	<i>Mesoplodon bowdoini</i>	Ziphiidae	1960-1969
				Antarctic Petrel	<i>Thalassica antarctica</i>	Procellariidae	1990-1999
				Antarctic Prion	<i>Pachyptila desolata</i>	Procellariidae	2000-2010
f				Apostlebird	<i>Struthidea cinerea</i>	Corcoracidae	2000-2010
				Arctic Jaeger	<i>Stercorarius parasiticus</i>	Laridae	2000-2010
				Arctic Tern	<i>Sterna paradisaea</i>	Laridae	2000-2010
				Asian Dowitcher	<i>Limnodromus semipalmatus</i>	Scolopacidae	1980-1989
f		e		Australasian Bittern	<i>Botaurus poiciloptilus</i>	Ardeidae	2000-2010
				Australasian Figbird	<i>Sphecotheres viridis</i>	Oriolidae	2000-2010
				Australasian Gannet	<i>Morus serrator</i>	Sulidae	2000-2010
				Australasian Grebe	<i>Tachybaptus novaehollandiae</i>	Pedicipedidae	2000-2010
				Australasian Pipit	<i>Anthus novaeseelandiae</i>	Motacillidae	2000-2010
				Australasian Shearwater	<i>Puffinus pacificus</i>	Procellariidae	2000-2010

Enter the name or an abbreviation then click 'Find Name'

This is a full list of all terrestrial vertebrates known for Victoria. As each name is highlighted a description and image will be displayed.

General Appearance: A flightless, marine bird, to 75 cm long, with short, flipper-like wings. Plumage is black on the back, head and wings, and white on breast and belly. Head is characterised by a clear, white ring around eyes.


Beak: Short and stout, dark red with black tip.

Legs: Very short, set well back on body, with pink, webbed feet.

Sexes: Similar.

Environment: Marine waters.

Notes: The only Victorian record is on the south-western coast in 1933.



Photographer: Bill Emison

Exit

Help

Set Filters - Full

Apply Filters

Set Filters - Quick

Show Current Filters

Print or Export List in Alphabetic Order

Print or Export List in Systematic Order

Plot Distribution Map

Print Picture Text and Map

Display Larger Picture

Highlight any technical word then right-click mouse to find its meaning

Image 1 of 1

Just a Minute Victorian Animals - 1:1,000,000 to 1:100,000

Enter Common Name

alp eger

Find Name

Sort on Scientific Name

Common Name Order : Full List : 780 species

Click on column buttons for interpretation of this list

F	A	S	O	Common Name	Scientific Name	Family Name	Last Date
				Adelie Penguin	Pygoscelis adeliae	Spheniscidae	No Records
				Agile Antechinus	Antechinus agilis	Dasyuridae	2000-2010
f		e		Alpine Bog Skink	Pseudemoia cryodroma	Scincidae	2000-2010
f		c		Alpine Egernia	Egernia guthega	Scincidae	2000-2010
f		e		Alpine She-oak Skink	Cyclodomorphus praealtus	Scincidae	2000-2010
f	V	c		Alpine Tree Frog	Litoria verreauxii alpina	Hylidae	2000-2010
f		c		Alpine Water Skink	Eulamprus kosciuskoi	Scincidae	2000-2010
				Andrew's Beaked Whale	Mesoplodon bowdoini	Ziphiidae	1960-1969
				Antarctic Petrel	Thalassoica antarctica	Procellariidae	1990-1999
				Antarctic Prion	Pachyptila desolata	Procellariidae	2000-2010
f				Apostlebird	Struthidea cinerea	Corcoracidae	2000-2010
				Arctic Jaeger	Stercorarius parasiticus	Laridae	2000-2010
				Arctic Tern	Sterna paradisaea	Laridae	2000-2010
				Asian Dowitcher	Limnodromus semipalmatus	Scolopacidae	1980-1989
f		e		Australasian Bittern	Botaurus poiciloptilus	Ardeidae	2000-2010
				Australasian Figbird	Sphecotheres viridis	Sulidae	2000-2010
				Australasian Gannet	Morus serrator	Pelecanidae	2000-2010
				Australasian Grebe	Tachybaptus novaehollandiae	Podicipedidae	2000-2010
				Australasian Pipit	Anthus novaeseelandiae	Motacillidae	2000-2010
				Australasian Shearwater	Puffinus pacificus	Procellariidae	2000-2010

General Appearance:

A relatively broad-headed, long-tailed, smooth-scaled lizard. Upper surface very variable, from pale brown to dark brown, with two broad, darker (often broken) stripes, from nape to hind legs, either side of the backbone, each containing variable pale brown spots. Flanks black, with pale brown to pale grey blotches. Head usually uniform in colour above and with dark brown stripes above and below the eye, and white or cream lips and eyelids. Lower surface off-white with a few darker markings. Combined length of legs is about the same as, or greater than, the distance between them. Feet have five toes; fourth toe on hind foot much longer than the third.

Snout-vent Length:

to 110 mm.

Tail:

to 140 mm.

Scalation:

Scales at mid-body, in 35-40 rows. Parietal scales not touching behind the interparietal scale. Interparietal scale much narrower than frontal-scale in adults.

Breeding:

Gives birth to live young.

Diet:

Principally arthropods. Diurnal.

Environment:

A rocky habitats, chiefly in high altitude areas.

Click here to display larger images of Alpine Egernia

Photographer: Nick Clemann/Viridans Images

Image 1 of 2

Exit

Help

Set Filters - Full

Apply Filters

Set Filters - Quick

Show Current Filters

Print or Export List in Alphabetic Order

Print or Export List in Systematic Order


Plot Distribution Map

Print Picture Text and Map

Display Larger Picture

Just a Minute Victorian Animals - 1:1,000,000 to 1:100,000

Alpine Egernia



Click here to view more images

Next

X

Click here to close image


family Name	Last Date
pheniscidae	No Records
asyuridae	2000-2010
cincidae	2000-2010
cincidae	2000-2010
cincidae	2000-2010
ylidae	2000-2010
cincidae	2000-2010
phiidae	1960-1969
ocellariidae	1990-1999
ocellariidae	2000-2010
orcoracidae	2000-2010
uridae	2000-2010
uridae	2000-2010
colopacidae	1980-1989
deidae	2000-2010
iolidae	2000-2010
ulidae	2000-2010
bdicipedidae	2000-2010
otacillidae	2000-2010
etidae	2000-2010

h

with

ave

le



Photographer: Nick Clemann/Viridans Images

Exit

Help

Set Filters - Full

Apply Filters

Set Filters - Quick

Show Current Filters

Print or Export List in Alphabetic Order

Print or Export List in Systematic Order

Plot Distribution Map

Print Picture Text and Map

Display Larger Picture

Image 1 of 2

© Nick Clemann/Viridans Images

Breeding: Gives birth to live young.

Diet: Principally arthropods. Diurnal.

Environment: A rocky habitats, chiefly in high altitude areas.

Highlight any technical word then right-click mouse to find its meaning

Just a Minute Victorian Animals - 1:1,000,000 to 1:100,000

Enter Common Name

alp eger

Find Name

Sort on Scientific Name

Common Name Order : Full List : 780 species

Click on column buttons for interpretation of this list

F	A	S	O	Common Name	Scientific Name	Family Name	Last Date
				Adelie Penguin	Pygoscelis adeliae	Spheniscidae	No Records
				Agile Antechinus	Antechinus agilis	Dasyuridae	2000-2010
f		e		Alpine Bog Skink	Pseudemoia cryodroma	Scincidae	2000-2010
f		c		Alpine Egernia	Egernia guthega	Scincidae	2000-2010
f		e		Alpine She-oak Skink	Cyclodomorphus praealtus	Scincidae	2000-2010
f	V	c		Alpine Tree Frog	Litoria verreauxii alpina	Hylidae	2000-2010
f		c		Alpine Water Skink	Eulamprus kosciuskoi	Scincidae	2000-2010
				Andrew's Beaked Whale	Mesoplodon bowdoini	Ziphiidae	1960-1969
				Antarctic Petrel	Thalassoica antarctica	Procellariidae	1990-1999
				Antarctic Prion	Pachyptila desolata	Procellariidae	2000-2010
f				Apostlebird	Struthidea cinerea	Corcoracidae	2000-2010
				Arctic Jaeger	Stercorarius parasiticus	Laridae	2000-2010
				Arctic Tern	Sterna paradisaea	Laridae	2000-2010
				Asian Dowitcher	Limnodromus semipalmatus	Scolopacidae	1980-1989
f		e		Australasian Bittern	Botaurus poiciloptilus	Ardeidae	2000-2010
				Australasian Figbird	Sphecotheres viridis	Ornithidae	2000-2010
				Australasian Gannet	Morus serrator	Sulidae	2000-2010
				Australasian Grebe	Tachybaptus novaehollandiae	Podicipedidae	2000-2010
				Australasian Pipit	Anthus novaeseelandiae	Motacillidae	2000-2010
				Australasian Shearwater	Puffinus pacificus	Procellariidae	2000-2010

General Appearance:

A relatively broad-headed, long-tailed, smooth-scaled lizard. Upper surface very variable, from pale brown to dark brown, with two broad, darker (often broken) stripes, from nape to hind legs, either side of the backbone, each containing variable pale brown spots. Flanks black, with pale brown to pale grey blotches. Head usually uniform in colour above and with dark brown stripes above and below the eye, and white or cream lips and eyelids. Lower surface off-white with a few darker markings. Combined length of legs is about the same as, or greater than, the distance between them. Feet have five toes; fourth toe on hind foot much longer than the third.

Snout-vent Length:

to 110 mm.

Tail:

to 140 mm.

Scalation:

Scales at mid-body, in 35-40 rows. Parietal scales not touching behind the interparietal scale. Interparietal scale much narrower than frontal-scale in adults.

Breeding:

Gives birth to live young.

Diet:

Principally arthropods. Diurnal.

Environment:

A rocky habitats, chiefly in high altitude areas.

Click here to display a summary page

Print Picture Text and Map

Display Larger

Create and print the selected image and text in an A4 format

Exit

Help

Set Filters - Full

Apply Filters

Set Filters - Quick

Show Current Filters

Print or Export List in Alphabetic Order

Print or Export List in Systematic Order

Plot Distribution Map

Photographer: Nick Clemann/Viridans Images

Image 1 of 2


Print

Printer
SetupReset
Picture

Reset Map

Help

Exit

Click here to send this
page to a printer**f c Alpine Egernia - *Egernia guthega****Photo by: Nick Clemann/Viridans Images*

General Appearance: A relatively broad-headed, long-tailed, smooth-scaled lizard. Upper surface very variable, from pale brown to dark brown, with two broad, darker (often broken) stripes, from nape to hind legs, either side of the backbone, each containing variable pale brown spots. Flanks black, with pale brown to pale grey blotches. Head usually uniform in colour above and with dark brown stripes above and below the eye, and white or cream lips and eyelids. Lower surface off-white with a few darker markings. Combined length of legs is about the same as, or greater than, the distance between them. Feet have five toes; fourth toe on hind foot much longer than the third.

Snout-vent Length: to 110 mm.

Tail: to 140 mm.

Scalation: Scales at mid-body, in 35-40 rows. Parietal scales not touching behind the interparietal scale. Interparietal scale much narrower than frontal-scale in adults.

Breeding: Gives birth to live young.

Diet: Principally arthropods. Diurnal.

Environment: A rocky habitats, chiefly in high altitude areas.

Notes: Until recently regarded as part of *Egernia whitii*. Differs from that species in its black flanks, more complex pattern on the back and by a proportionately shorter tail.

Conservation: **ffl** Listed under the Victorian Flora and Fauna Guarantee. **[c]** Critically Endangered in Victoria.

Just a Minute Victorian Animals - 1:1,000,000 to 1:100,000

Enter Common Name

alp eger

Find Name

Sort on Scientific Name

Common Name Order : Full List : 780 species

Click on column buttons for interpretation of this list

F	A	S	O	Common Name	Scientific Name	Family Name	Last Date
				Adelie Penguin	Pygoscelis adeliae	Spheniscidae	No Records
				Agile Antechinus	Antechinus agilis	Dasyuridae	2000-2010
f		e		Alpine Bog Skink	Pseudemoia cryodroma	Scincidae	2000-2010
f		c		Alpine Egernia	Egernia guthega	Scincidae	2000-2010
f		e		Alpine She-oak Skink	Cyclodomorphus praealtus	Scincidae	2000-2010
f	V	c		Alpine Tree Frog	Litoria verreauxii alpina	Hylidae	2000-2010
f		c		Alpine Water Skink	Eulamprus kosciuskoi	Scincidae	2000-2010
				Andrew's Beaked Whale	Mesoplodon bowdoini	Ziphiidae	1960-1969
				Antarctic Petrel	Thalassoica antarctica	Procellariidae	1990-1999
				Antarctic Prion	Pachyptila desolata	Procellariidae	2000-2010
f				Apostlebird	Struthidea cinerea	Corcoracidae	2000-2010
				Arctic Jaeger	Stercorarius parasiticus	Laridae	2000-2010
				Arctic Tern	Sterna paradisaea	Laridae	2000-2010
				Asian Dowitcher	Limnodromus semipalmatus	Scolopacidae	1980-1989
f		e		Australasian Bittern	Botaurus poiciloptilus	Ardeidae	2000-2010
				Australasian Figbird	Sphecotheres viridis	Oriolidae	2000-2010
				Australasian Gannet	Morus serrator	Sulidae	2000-2010
				Australasian Grebe	Tachybaptus novaehollandiae	Podicipedidae	2000-2010
				Australasian Pipit	Anthus novaeseelandiae	Motacillidae	2000-2010
				Australasian Shearwater	Puffinus pacificus	Procellariidae	2000-2010

Click here to generate a Distribution Map

Plot distribution of highlighted species on a map

Print or Export List in Alphabetic Order

Print or Export List in Systematic Order

Plot Distribution Map

General Appearance:

A relatively broad-headed, long-tailed, smooth-scaled lizard. Upper surface very variable, from pale brown to dark brown, with two broad, darker (often broken) stripes, from nape to hind legs, either side of the backbone, each containing variable pale brown spots. Flanks black, with pale brown to pale grey blotches. Head usually uniform in colour above and with dark brown stripes above and below the eye, and white or cream lips and eyelids. Lower surface off-white with a few darker markings. Combined length of legs is about the same as, or greater than, the distance between them. Feet have five toes; fourth toe on hind foot much longer than the third.

Snout-vent Length:

to 110 mm.

Tail:

to 140 mm.

Scalation:

Scales at mid-body, in 35-40 rows. Parietal scales not touching behind the interparietal scale. Interparietal scale much narrower than frontal-scale in adults.

Breeding:

Gives birth to live young.

Diet:

Principally arthropods. Diurnal.

Environment:

A rocky habitats, chiefly in high altitude areas.

Photographer: Nick Clemann/Viridans Images

Print Picture Text and Map

Display Larger Picture

Image 1 of 2

Highlight any technical word then right-click mouse to find its meaning

