

Viridans FIS-VFD Tutorial 2 - Web

Task:

Use the VFD to find the rare or threatened animals of the Ouyen area.

- Locate Ouyen on a map of Victoria.
- Draw a circle of 10 km radius centred on Ouyen.
- Search the VFD database for all records within that area.
- Save a map of this search in a PDF file.
- Create a list of all species within the search area.
- Generate a subset of that list containing only species categorised as rare or threatened.
- Export the list as a PDF file that can be loaded into a word processor.

The products of this task should be:

- A PDF file containing a list of rare and threatened animals found within 10 km of Ouyen.
- A map, in PDF format, of part of Victoria with all sites from the search area superimposed.

Find a Species

Search an Area

Access Site Data

General Information

FAQ

Logout

Victorian
Fauna Database



2013

Copyright © Viridans Biological Databases

Back



Victorian Fauna Database

Search from Base Map
Use regional map with point-and-click to search [Click here to find place](#)

Search using Place Name
Enter Place Name (minimum 3 characters)
ouyen [Enter Place Name](#)

Search using Geocode

Longitude (DMS)		Latitude (DMS)	
<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
7 digits		6 digits	
Zone	Easting	Northing	
<input type="text"/>	<input type="text"/>	<input type="text"/>	
2 digits	6 digits	7 digits	

Define a Circular Search Area

- No pre-defined search area
- No pre-defined search area
- Circle Radius 3 km
- Circle Radius 4 km
- Circle Radius 5 km
- Circle Radius 6 km
- Circle Radius 7 km
- Circle Radius 8 km
- Circle Radius 9 km
- Circle Radius 10 km**
- Circle Radius 11 km
- Circle Radius 12 km

[Select Search area radius](#)

[Back](#)




Victorian Fauna Database

Places beginning with - Ouyen

Ouyen plus 10 km radius	Township	142°18'43" 35°04'16"	
Ouyen Channel plus 10 km radius	Place Name	142°31'31" 35°09'21"	
Ouyen I34 B.R. plus 10 km radius	Bushland Reseve	142°19'36" 34°59'50"	
Ouyen I82 B.R. plus 10 km radius	Bushland Reseve	142°18'17" 35°04'32"	
Ouyen I83 B.R. plus 10 km radius	Bushland Reseve	142°21'50" 35°03'49"	
Ouyen I85 B.R. plus 10 km radius	Bushland Reseve	142°22'31" 35°05'57"	
Ouyen Magistrates Court plus 10 km radius	Place Name	142°18'55" 35°04'09"	

Click here to print or export the map





- 5 km
- Main Roads
 - Sealed Roads
 - Unsealed Roads
 - Tracks
 - Built Up Areas
 - Urban Parks
 - Sea
 - Rivers
 - Lakes
 - Reservoirs
 - Swamps
 - Private Land
 - Public Land
 - National Parks
 - State Parks
 - Regional Parks
 - Wildlife Reserves
 - Wilderness Areas
 - Reference Areas



- Map Options*
- Sketch a Search Area
 - Search Sketched Area
 - Get all data from sites
 - Identify sites on map
 - Map Zoom

Data modified from Viridans Victorian Fauna Database

```
javascript: findallsites1();
```



Click anywhere on the map to zoom



5 km

- Main Roads
- Sealed Roads
- Unsealed Roads
- Tracks
- Built Up Areas
- Urban Parks
- Sea
- Rivers
- Lakes
- Reservoirs
- Swamps
- Private Land
- Public Land
- National Parks
- State Parks
- Regional Parks
- Wildlife Reserves
- Wilderness Areas
- Reference Areas

Click here to summarise data from all sites

- Map Options*
- Sketch a Search Area
 - Search Sketched Area
 - Get all data from sites
 - Identify sites on map
 - Map Zoom

Data modified from Viridans Victorian Fauna Database

```
javascript: findallsites1();
```

Close

Show Filter Options

List Title: Rare Animals of Ouyen Area

Print-Save PDF Save CSV

Acanthagenys rufogularis

Photo: © Wendy Opie/Viridans Images

General Appearance: A brown and yellow bird, to 25 cm long. Plumage predominantly dark grey-brown, streaked with pale brown, on head, back and wings. Breast yellow. Belly pale yellow to pale grey with dark brown spots. Head characterised by a dark pink eye ring and a broad band of bristly white feathers beneath the eye, running from beak to cheeks (hence the common name).

Tail: About as long as body. Dark grey-brown with white tips.

Eyes: Blue.

Beak: Narrow, curved downwards. Pink with a black tip.

Legs: Dark grey.

Sexes: Similar.

Call: A melodic warbling and a repeated trill.

Juveniles: Similar to adults but duller.

Eggs: Off-white sometimes with a few darker spots (Jul - Jan).

Nest: A sparse cup made from grass and fine twigs.



Click here to display filter options

Plot Distribution Map Show Larger Pictures Display Picture Text and Map

A B C D E F G H I J K L M N O P Q R S T U V W X Y Z

F A V O Species Names	Family	Records
<i>Acanthagenys rufogularis</i> - Spiny-cheeked Honeyeater	Meliphagidae	14 : 5.2%
<i>Acanthiza chrysorrhoa</i> - Yellow-rumped Thornbill	Pardalotidae	15 : 5.57%
<i>Acanthiza nana</i> - Yellow Thornbill	Pardalotidae	1 : 0.37%
<i>Acanthiza reguloides</i> - Buff-rumped Thornbill	Pardalotidae	1 : 0.37%
<i>Acanthiza uropygialis</i> - Chestnut-rumped Thornbill	Pardalotidae	8 : 2.97%
<i>Accipiter fasciatus</i> - Brown Goshawk	Accipitridae	1 : 0.37%
<i>Acrocephalus stentoreus</i> - Clamorous Reed Warbler	Sylviidae	1 : 0.37%
<i>Amphibolurus nobbi coggeri</i> - Nobbi Dragon	Agamidae	9 : 3.34%
<i>Anas gracilis</i> - Grey Teal	Anatidae	1 : 0.37%
<i>Anas superciliosa</i> - Pacific Black Duck	Anatidae	2 : 0.74%
<i>Anthochaera carunculata</i> - Red Wattlebird	Meliphagidae	15 : 5.57%
<i>Anthus novaeseelandiae</i> - Australasian Pipit	Motacillidae	5 : 1.85%
<i>Aphelocephala leucopsis</i> - Southern Whiteface	Pardalotidae	4 : 1.48%
<i>Aprasia aurita</i> - Mallee Worm-lizard	Pygopodidae	12 : 4.46%
<i>Aprasia inaurita</i> - Pink-nosed Worm-Lizard	Pygopodidae	2 : 0.74%
<i>Aprasia sp.</i> - Aprasia	Pygopodidae	1 : 0.37%
<i>Aquila audax</i> - Wedge-tailed Eagle	Accipitridae	3 : 1.11%
<i>Artamus cinereus</i> - Black-faced Woodswallow	Artamidae	2 : 0.74%
<i>Artamus personatus</i> - Masked Woodswallow	Artamidae	2 : 0.74%
<i>Artamus superciliosus</i> - White-browed Woodswallow	Artamidae	17 : 6.31%

This list contains all animal species recorded from the search area.

135 Species

Copyright © Viridans Biological Databases

Refine this List Reset all Filters

Display Order Conservation Animal Group Breeding Seasons Period of Activity Diet

Scientific Name Order All Categories All Animal Groups Any Season Any Time All Types

Acanthagenys rufogularis

General Appearance: Acanthagenys rufogularis is a small bird, approximately 10 cm long. Plumage predominantly dark grey-brown, streaked with pale brown. Rump pale grey-brown. Throat and breast yellow. Belly pale yellow to pale grey with dark streaks. Characterised by a dark pink eye ring and a broad band of dark to black to cheeks (hence the common name).

Tail: About as long as body.

Eyes: Blue.

Beak: Narrow, curved downwards.

Legs: Dark grey.

Sexes: Similar.

Call: A melodic warbling and a repeated trill.

Juveniles: Similar to adults but duller.

Eggs: Off-white sometimes with a few darker spots (Jul - Jan).

Nest: A sparse cup made from grass and fine twigs.

- All Categories
- Just Native Species
- Just Non-native Species
- Rare or threatened species
- Listed under EPBC
- Listed under FFG
- CAMBA Treaty
- JAMBA Treaty
- Hollow Dependence
- Migrants
- Vagrants
- Residents

Select the Rare or Threatened option



Plot Distribution Map Show Larger Pictures Display Picture Text and Map

A B C D E F G H I J K L M N O P Q R S T U V W X Y Z

F A V O Species Names	Family	Records
<i>Acanthagenys rufogularis</i> - Spiny-cheeked Honeyeater	Meliphagidae	14 : 5.2%
<i>Acanthiza chrysorrhoa</i> - Yellow-rumped Thornbill	Pardalotidae	15 : 5.57%
<i>Acanthiza nana</i> - Yellow Thornbill	Pardalotidae	1 : 0.37%
<i>Acanthiza reguloides</i> - Buff-rumped Thornbill	Pardalotidae	1 : 0.37%
<i>Acanthiza uropygialis</i> - Chestnut-rumped Thornbill	Pardalotidae	8 : 2.97%
<i>Accipiter fasciatus</i> - Brown Goshawk	Accipitridae	1 : 0.37%
<i>Acrocephalus stentoreus</i> - Clamorous Reed Warbler	Sylviidae	1 : 0.37%
<i>Amphibolurus nobbi coggeri</i> - Nobbi Dragon	Agamidae	9 : 3.34%
<i>Anas gracilis</i> - Grey Teal	Anatidae	1 : 0.37%
<i>Anas superciliosa</i> - Pacific Black Duck	Anatidae	2 : 0.74%
<i>Anthochaera carunculata</i> - Red Wattlebird	Meliphagidae	15 : 5.57%
<i>Anthus novaeseelandiae</i> - Australasian Pipit	Motacillidae	5 : 1.85%
<i>Aphelocephala leucopsis</i> - Southern Whiteface	Pardalotidae	4 : 1.48%
<i>Aprasia aurita</i> - Mallee Worm-lizard	Pygopodidae	12 : 4.46%
<i>Aprasia inaurita</i> - Pink-nosed Worm-Lizard	Pygopodidae	2 : 0.74%
<i>Aprasia sp.</i> - Aprasia	Pygopodidae	1 : 0.37%
<i>Aquila audax</i> - Wedge-tailed Eagle	Accipitridae	3 : 1.11%
<i>Artamus leucorhynchus</i> - Black-faced Woodswallow	Artamidae	2 : 0.74%

[Close](#) [Hide Filter Options](#) List Title: **Rare Animals of Ouyen Area** [Print-Save PDF](#) [Save CSV](#)

[Refine this List](#) [Reset all Filters](#)

Display Order: Scientific Name Order
 Conservation: **Rare or threatened species**
 Animal Group: All Animal Groups
 Breeding Seasons: Any Season
 Period of Activity: Any Time
 Diet: All Types

Click here to filter the list

Acanthagenys rufogularis

General Appearance: A brown and yellow bird, to 25 cm long. Plumage predominantly dark grey-brown, streaked with pale brown, on head, back and upper wings. Rump pale grey-brown. Throat and breast yellow. Belly pale yellow to pale grey with dark brown spots. Head characterised by a dark pink eye ring and a broad band of bristly white feathers beneath the eye, running from beak to cheeks (hence the common name).

Tail: About as long as body. Dark grey-brown with white tips.

Eyes: Blue.

Beak: Narrow, curved downwards. Pink with a black tip.

Legs: Dark grey.

Sexes: Similar.

Call: A melodic warbling and a repeated trill.

Juveniles: Similar to adults but duller.

Eggs: Off-white sometimes with a few darker spots (Jul - Jan).

Nest: A sparse cup made from grass and fine twigs.



[Plot Distribution Map](#) [Show Larger Pictures](#) [Display Picture Text and Map](#)

[A](#) [B](#) [C](#) [D](#) [E](#) [F](#) [G](#) [H](#) [I](#) [J](#) [K](#) [L](#) [M](#) [N](#) [O](#) [P](#) [Q](#) [R](#) [S](#) [T](#) [U](#) [V](#) [W](#) [X](#) [Y](#) [Z](#)

F A V O Species Names	Family	Records
Acanthagenys rufogularis - Spiny-cheeked Honeyeater	Meliphagidae	14 : 5.2%
Acanthiza chrysorrhoa - Yellow-rumped Thornbill	Pardalotidae	15 : 5.57%
Acanthiza nana - Yellow Thornbill	Pardalotidae	1 : 0.37%
Acanthiza reguloides - Buff-rumped Thornbill	Pardalotidae	1 : 0.37%
Acanthiza uropygialis - Chestnut-rumped Thornbill	Pardalotidae	8 : 2.97%
Accipiter fasciatus - Brown Goshawk	Accipitridae	1 : 0.37%
Acrocephalus stentoreus - Clamorous Reed Warbler	Sylviidae	1 : 0.37%
Amphibolurus nobbi coggeri - Nobbi Dragon	Agamidae	9 : 3.34%
Anas gracilis - Grey Teal	Anatidae	1 : 0.37%
Anas superciliosa - Pacific Black Duck	Anatidae	2 : 0.74%
Anthochaera carunculata - Red Wattlebird	Meliphagidae	15 : 5.57%
Anthus novaeseelandiae - Australasian Pipit	Motacillidae	5 : 1.85%
Aphelocephala leucopsis - Southern Whiteface	Pardalotidae	4 : 1.48%
Aprasia aurita - Mallee Worm-lizard	Pygopodidae	12 : 4.46%
Aprasia inaurita - Pink-nosed Worm-Lizard	Pygopodidae	2 : 0.74%
Aprasia sp. - Aprasia	Pygopodidae	1 : 0.37%
Aquila audax - Wedge-tailed Eagle	Accipitridae	3 : 1.11%
Artamus leucorhynchus - Black-faced Woodswallow	Artamidae	2 : 0.74%

Close

Show Filter Options

List Title: Rare Animals of Ouyen Area

Print-Save PDF

Save CSV

Acanthagenys rufogularis

Photo: © Wendy Opie/Viridans Images

General Appearance: A brown and yellow bird, to 25 cm long. Plumage predominantly dark grey-brown, streaked with pale brown, on head, back and upper wings. Underparts grey-brown. Breast yellow. Belly pale yellow to pale grey with dark brown spots. Head characterised by a grey eye ring and a broad band of bristly white feathers beneath the eye, running from beak to cheeks (hence the common name).

Tail: About as long as body. Dark grey-brown with white tips.

Eyes: Blue.

Beak: Narrow, curved downwards. Pink with a black tip.

Legs: Dark grey.

Sexes: Similar.

Call: A melodic warbling and a repeated trill.

Juveniles: Similar to adults but duller.

Eggs: Off-white sometimes with a few darker spots (Jul - Jan).

Nest: A sparse cup made from grass and fine twigs.

Change the title from the default

Click here to export the list in PDF format



Plot Distribution Map Show Larger Pictures Display Picture Text and Map

A B C D E F G H I J K L M N O P Q R S T U V W X Y Z

F A V O Species Names Family Records

n	<i>Aprasia aurita</i> - Mallee Worm-lizard	Pygopodidae	12 : 4.46%
n	<i>Cercartetus concinnus</i> - Western Pygmy-possum	Burramyidae	1 : 0.37%
n	<i>Climacteris picumnus victoriae</i> - Brown Treecreeper (south-eastern esp.)	Climacteridae	6 : 2.23%
v	<i>Falco subniger</i> - Black Falcon	Falconidae	1 : 0.37%
f	<i>Leoprocra leadbeateri</i> - Major Mitchell's Cockatoo	Cacatuidae	2 : 0.74%
f	<i>Melanodryas cucullata</i> - Hooded Robin	Petroicidae	1 : 0.37%
f	<i>Polytelis anthopeplus</i> - Regent Parrot	Psittacidae	5 : 1.85%
d	<i>Pseudonaja nuchalis</i> - Western Brown Snake	Elapidae	1 : 0.37%
d	<i>Tiliqua occipitalis</i> - Western Blue-tongued Lizard	Scincidae	1 : 0.37%
f	<i>Vermicella annulata</i> - Bandy Bandy	Elapidae	2 : 0.74%

The list contains all rare or threatened species from the search area

Victorian rare or threatened
Listed under the EPBC Act
Listed under the FFG Act

10 Species

Copyright © Viridans Biological Databases

javascript: alphaspec("speclistgrid1.pdf.php");

localhost/viridans/vfddesk/specistgrid1.pdf.php?namestatus=c...

localhost/viridans/vfddesk/specistgrid1.pdf.php?namestatus=cursci&conserv=1

Rare Animals of Ouyen Area

f n	<i>Aprasia acrita</i>	Mallee Worm-lizard	Pygopodidae	12 : 4.46%
n	<i>Dorcaterus concoloratus</i>	Western Pygmy-possum	Burramyidae	1 : 0.37%
n	<i>Climacteris picavetius vicitorius</i>	Brown Treecreeper (south-eastern ssp.)	Climacteridae	6 : 2.23%
v	<i>Falco subniger</i>	Black Falcon	Falconidae	1 : 0.37%
f v	<i>Lophoscroa leachbeateri</i>	Major Mitchell's Cockatoo	Cacatuidae	2 : 0.74%
f n	<i>Melanodryas cucullata</i>	Hooded Robin	Petroicidae	1 : 0.37%
f v v	<i>Polytelis anthrapapulus</i>	Regent Parrot	Psittacidae	5 : 1.85%
d	<i>Pseudonaja nuchalis</i>	Western Brown Snake	Elapidae	1 : 0.37%
d	<i>Tiliqua occipitalis</i>	Western Blue-tongued Lizard	Scincidae	1 : 0.37%
f n	<i>Vernicellia annulata</i>	Bandy Bandy	Elapidae	2 : 0.74%

The PDF file can be saved, printed or emailed.

Control

10 Species

List Title: **Rare Animals of Ouyen Area**

Print-Save PDF Save CSV

Photo: © Wendy Opie/Viridans Images

25 cm long. Plumage predominantly dark grey-brown, wings. Rump pale grey-brown. Throat and breast yellow. Belly and underparts characterised by a dark pink eye ring and a broad band of white tips.



tip.

Jul - Jan).

Show Larger Pictures

Display Picture Text and Map

I J K L M N O P Q R S T U V W X Y Z

	Family	Records
-possum	Pygopodidae	12 : 4.46%
Treecreeper (south-eastern ssp.)	Burramyidae	1 : 0.37%
Cockatoo	Climacteridae	6 : 2.23%
	Falconidae	1 : 0.37%
	Cacatuidae	2 : 0.74%
	Petroicidae	1 : 0.37%
Snake	Psittacidae	5 : 1.85%
Red Lizard	Elapidae	1 : 0.37%
	Scincidae	1 : 0.37%
	Elapidae	2 : 0.74%

Copyright © Viridans Biological Databases